



Uniqueness result for an age-dependent reaction-diffusion problem

Vo Anh Khoa, Tran The Hung, Daniel Lesnic

UHasselT Computational Mathematics Preprint
Nr. UP-18-07

Oct. 2, 2018

UNIQUENESS RESULT FOR AN AGE-DEPENDENT REACTION-DIFFUSION PROBLEM

VO ANH KHOA, TRAN THE HUNG, AND DANIEL LESNIC

ABSTRACT. This paper is concerned with an age-structured model in population dynamics. We investigate the uniqueness of solution for this type of nonlinear reaction-diffusion problem when the source term depends on the density, indicating the presence of, for example, mortality and reaction processes. Our result shows that in a spatial environment, if two population densities obey the same evolution equation and possess the same terminal data of time and age, then their distributions must coincide therein.

1. INTRODUCTION

In population dynamics, there are several factors interestingly contributing to the complicated and nontrivial spatio-temporal spread patterns of diseases. Especially demographic-disease ages and spatial movement naturally interweave in many ways. Their correlation and interaction are expected to participate in mathematical modeling and analysis, and can lead to models with distinct levels of complexity.

As far as we know, disease management measures are often age-dependent and their effectiveness may also be dependent on the mobility status of the involved species, such as larvicides and insecticide sprays for mosquito-borne diseases (e.g. the invasion and spread of West Nile virus in North America in [8]) used to control the mosquito population at different levels of their maturity. We recall the structured population model in [12] for the total number of matured individuals in a single species population at the demographic age, which reads as

$$(1.1) \quad u_t + u_a = D\Delta u - d_0 u.$$

Here, t denotes the time, a denotes age, u represents the population density, D and d_0 are, respectively, the diffusion and death rates of the immature individual, under the assumption that the maturation rate is regulated by the birth process and the dynamics of the individual during the maturation phase.

A generalized approach for this mosquito-borne disease model is given by the coupled McKendrick von-Foerster equations (see also [11, 13, 14]) where the reservoir as the host and the age-structured host population are observed. Following [2], the spatial spread of vector-borne diseases model involving age-structure is

$$(1.2) \quad \begin{cases} u_t + u_a = D_1(a) \Delta u - u(d_1(a) + \chi(a) m(t, x)), \\ v_t + v_a = D_2(a) \Delta v - v d_2(a) + u \chi(a) m(t, x). \end{cases}$$

Here, u and v play roles as susceptible and infected hosts, respectively, while $D_i(a)$ are the age-dependent diffusivities by natural means, $\chi(a)$ is the age-dependent

2000 *Mathematics Subject Classification.* 35A07, 35K57, 92D25, 45Q05.

Key words and phrases. Backward age-dependent reaction-diffusion, Uniqueness, Population dynamics.

transmission coefficient, $d_i(a)$ are the age-dependent death rates of such hosts for $i = 1, 2$ and m can be regarded as the number of infected adult mosquitoes.

Let $\Omega \subset \mathbb{R}^n$ for $n \geq 2$ be a bounded, open and connected domain with sufficiently smooth boundary and let \mathcal{A} be the linear second-order differential operator

$$(1.3) \quad \mathcal{A}u(t, a, x) = \sum_{i,j=1}^n \frac{\partial}{\partial x_i} \left(d_{ij}(t, a, x) \frac{\partial u(t, a, x)}{\partial x_j} \right),$$

accounting e.g. for the anisotropic diffusion and possibly taxis processes.

Setting $Q_{T,a_\dagger}^\Omega := (0, T) \times (0, a_\dagger) \times \Omega$ for $T, a_\dagger > 0$, we consider the problem:

$$(1.4) \quad \begin{cases} u_t + u_a - \mathcal{A}u = F(t, a, x; u) & \text{in } Q_{T,a_\dagger}^\Omega, \\ u(T, a, x) = u_0(a, x) & \text{in } (0, a_\dagger) \times \Omega, \\ u(t, a_\dagger, x) = u_1(t, x) & \text{in } (0, T) \times \Omega, \end{cases}$$

where F is a given source term, endowed either with the homogeneous Dirichlet boundary condition

$$(1.5) \quad u(t, a, x) = 0 \quad \text{on } (0, T) \times (0, a_\dagger) \times \partial\Omega,$$

or with the Robin-type boundary condition

$$(1.6) \quad -d(t, a, x) \nabla u(t, a, x) \cdot \mathbf{n} = S(u) \quad \text{on } (0, T) \times (0, a_\dagger) \times \partial\Omega.$$

Here, \mathbf{n} denotes the outward unit normal vector to the boundary $\partial\Omega$, $d = (d_{ij})_{i,j=1,\dots,n}$ is the diffusion tensor and S is some given function.

In this paper, we prove that under certain assumptions on the source term F and surface reaction S , the solution to the system (1.4) and (1.5) or (1.6) is unique, if it exists. The present source term F can model several types of polynomial reactions (e.g. logistic and von Bertalanffy) and even exponential growth (Arrhenius laws). It can also be modelled as

$$F(t, a, x; u) = -\tilde{\mu}(t; u) u(t, a, x),$$

where $\tilde{\mu} > 0$ is called the time- and density-dependent mortality modulus. This mortality-related functional usually arises in the Lotka-von Foerster model, where the simple modulus is

$$\tilde{\mu}(t; u) = \int_0^{a_\dagger} \int_\Omega u(t, a, x) dx da,$$

provided the mortality process is also controlled by the total population at time t during the whole age and environment.

It is worth noting that when age can be viewed as a temporal time, the first equation in (1.4) is usually referred to as an ultra-parabolic equation. In literature, the backward problem (1.4) and (1.5) has been explored in [15] for an abstract linear class of ultra-parabolic problems. Such problems are basically ill-posed in the sense that the solution does not depend continuously on the data no matter how smooth it is. As a matter of fact, a regularization has to be designed. The starting point of an evolution equation involving multi-time variables is from [6, 7]. It is shown in [6] that the forward problem for (1.4) and (1.5) is well-posed in the space of Hölder continuous functions. As a byproduct, the result therein provides the representation of solution, based on the semi-group theory along the upper and lower triangles dividing the rectangle of times. The evolution of the system with two different times were discussed by an argument where some diffusion processes

with memory take place. In the framework of stochastic optimal control, the reader can be referred to e.g. [9, 10], for related contributions to this problem.

The rest of this paper is organized as follows: Section 2 is devoted to introduce notation and conventions throughout this paper. In addition, we provide technical assumptions on parameters and coefficients involved in (1.4). In Section 3, we deliberately present two subsections where the Dirichlet and Robin-type boundary conditions are considered, respectively. Our main results are reported in Theorem 3.2 and Theorem 3.5, whilst their cores of proof are based on Lemma 3.1 and Lemma 3.4, respectively. Essentially, the main technique here is inspired from [1, Chapter 6]. This approach was used to treat the backward parabolic operator within the study of the large-time behavior of solutions to a linear class of initial-boundary value parabolic equations. It is also helpful in the analysis of regularization for inverse and ill-posed problems. In this work, we extend the uniqueness result particularly to a class of semi-linear age-dependent reaction-diffusion equations.

2. PRELIMINARIES

In parallel with using the notation Q_{T,a_\dagger}^Ω , the same meaning is also given to the notation $Q_{T,a_\dagger} = (0, T) \times (0, a_\dagger)$ and $Q_{(t_1, t_2) \times (a_1, a_2)}^\Omega = (t_1, t_2) \times (a_1, a_2) \times \Omega$, etc. throughout this paper. Moreover, depending on the context, by $|\cdot|$ we denote either the volume measure of a domain or the absolute value of a function.

To this end, for any domain D we also set $\|\cdot\|_D$ to be the norm in $L^2(D)$ and the same indication goes to the inner product $\langle \cdot, \cdot \rangle_D$.

For the sake of our analysis in this work, we need the following set of assumptions:

(A₁) The diffusion tensor $d = (d_{ij})_{1 \leq i, j \leq n} \in [C^1(\overline{Q_{T,a_\dagger}^\Omega})]^{n \times n}$ is symmetric and there exist positive constants \underline{c} and \bar{c} such that

$$\underline{c}|\xi|^2 \leq \sum_{i,j=1}^n d_{ij}(t, a, x) \xi_i \xi_j \leq \bar{c}|\xi|^2 \quad \text{for any } \xi \in \mathbb{R}^n.$$

(A₂) There exists a positive constant $\bar{M} > 0$ such that

$$|\partial_t d_{ij}(t, a, x)| + |\partial_a d_{ij}(t, a, x)| \leq \bar{M} \quad \text{for all } (t, a, x) \in \overline{Q_{T,a_\dagger}^\Omega}.$$

(A₃) The source term $F : \overline{Q_{T,a_\dagger}^\Omega} \times L^2(Q_{T,a_\dagger}^\Omega) \rightarrow \mathbb{R}$ is $(0, 1] \ni \alpha$ -Hölder continuous with respect to the density argument, i.e. there exists $L_F > 0$ such that for all $(t, a, x) \in \overline{Q_{T,a_\dagger}^\Omega}$,

$$|F(t, a, x; u_1) - F(t, a, x; u_2)| \leq L_F |u_1 - u_2|^\alpha \quad \text{for all } u_1, u_2 \in L^2(Q_{T,a_\dagger}^\Omega).$$

(A₄) For the surface reaction term $S : L^2(\partial\Omega) \rightarrow \mathbb{R}$ there exists $\bar{m} > 0$ such that $\langle (\partial_t + \partial_a)(S(u_1) - S(u_2)), u_1 - u_2 \rangle_{\partial\Omega} \leq \bar{m} \|u_1 - u_2\|_{\partial\Omega}^2$ for all $u_1, u_2 \in L^2(\partial\Omega)$.

(A₅) The surface reaction term $S : L^2(\partial\Omega) \rightarrow \mathbb{R}$ is monotone, i.e.

$$\langle S(u_1) - S(u_2), u_1 - u_2 \rangle_{\partial\Omega} \geq 0 \quad \text{for all } u_1, u_2 \in L^2(\partial\Omega).$$

(A₆) The surface reaction term $S : L^2(\partial\Omega) \rightarrow \mathbb{R}$ is $(0, 1] \ni \beta$ -Hölder continuous, i.e. there exists $L_S > 0$ such that

$$|S(u_1) - S(u_2)| \leq L_S |u_1 - u_2|^\beta \quad \text{for all } u_1, u_2 \in L^2(\partial\Omega).$$

Lemma 2.1. (cf. [4]) For any $\gamma > 0$ and $\alpha_0 \in (0, 1]$, the following inequality holds

$$(2.1) \quad X^{\alpha_0} \leq \alpha_0 \gamma^{\alpha_0 - 1} X + (1 - \alpha_0) \gamma^{\alpha_0} \quad \text{for all } X \geq 0.$$

Lemma 2.2. (cf. [3]) There exists a positive constant C_0 such that

$$(2.2) \quad \|u\|_{\partial\Omega}^2 \leq C_0 \left(\|u\|_{\Omega}^2 + \|\nabla u\|_{\Omega}^2 \right) \quad \text{for any } u \in H^1(\Omega).$$

Remark 2.3. The constant C_0 in (2.2) can be identified from the trace inequality

$$(2.3) \quad \|u\|_{\partial\Omega}^2 \leq c(\varepsilon, \Omega) \|u\|_{\Omega}^2 + \varepsilon \|\nabla u\|_{\Omega}^2 \quad \text{for any } u \in H^1(\Omega) \text{ and } \varepsilon > 0.$$

This inequality is roughly dependent on the geometry of Ω . As a typical example from [3, Theorem 2], if Ω is star-shaped, $c(\varepsilon, \Omega)$ is of the order of $\mathcal{O}(1 + \varepsilon^{-1})$. When u has zero mean on $\partial\Omega$, i.e. $\int_{\partial\Omega} u d\sigma_x = 0$, then we have the stronger Poincaré inequality $\|u\|_{\partial\Omega}^2 \leq \tilde{C}_0 \|\nabla u\|_{\Omega}^2$, for some constant $\tilde{C}_0 > 0$.

3. MAIN RESULTS

3.1. Dirichlet boundary condition (1.5). Let us set the function space

$$(3.1) \quad W_{T, a_{\dagger}}^{\Omega} := C(\overline{Q_{T, a_{\dagger}}}) \cap H_0^1(\Omega) \cap L^2(Q_{T, a_{\dagger}}; H^2(\Omega)) \cap C^1(Q_{T, a_{\dagger}}; H_0^1(\Omega)).$$

Assume that there exist two solutions u_1 and u_2 of (1.4) and (1.5) that belong to $W_{T, a_{\dagger}}^{\Omega}$. Then their difference $w = u_1 - u_2 \in W_{T, a_{\dagger}}^{\Omega}$ satisfies

$$(3.2) \quad \begin{cases} w_t + w_a - \mathcal{A}w = F(t, a, x; u_1) - F(t, a, x; u_2) & \text{in } Q_{T, a_{\dagger}}^{\Omega}, \\ w(t, a, x) = 0 & \text{on } Q_{T, a_{\dagger}}^{\partial\Omega}, \\ w(T, a, x) = 0 & \text{in } (0, a_{\dagger}) \times \Omega, \\ w(t, a_{\dagger}, x) = 0 & \text{in } (0, T) \times \Omega. \end{cases}$$

Let us denote

$$P_{T, a_{\dagger}}^{\Omega} := \left\{ u \in W_{T, a_{\dagger}}^{\Omega} : u|_{\partial\Omega} = 0, u|_{t \in \{0, T\}} = 0, u|_{a \in \{0, a_{\dagger}\}} = 0 \right\}$$

and $\lambda(t, a) := t - T + a - a_{\dagger} - \eta < 0$ for some $\eta > 0$.

Lemma 3.1. Assume (A_1) and (A_2) hold. Then for any $v \in P_{T, a_{\dagger}}^{\Omega}$, $m \in \mathbb{N}^*$ and $k \in \mathbb{R}_+^*$, we have

$$(3.3) \quad \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}v - v_t - v_a) \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 \geq \frac{4m}{k} \left\| \lambda^{-\frac{m}{k} - 1} v \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 - \overline{M} \left\| \lambda^{-\frac{m}{k}} \nabla v \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2.$$

Moreover, if $0 < T + a_{\dagger} \leq \mu$ for a sufficiently small constant $\mu > 0$, there exists a positive K such that

$$(3.4) \quad K \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}v - v_t - v_a) \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 \geq \left\| \lambda^{-\frac{m}{k} - 1} v \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 + \frac{1}{2} \left\| \lambda^{-\frac{m}{k}} \nabla v \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2,$$

for sufficiently large m .

Proof. Let $z = \lambda^{-\frac{m}{k}} v \in P_{T, a_{\dagger}}^{\Omega}$. Then

$$(3.5) \quad \lambda^{-\frac{m}{k}} (\mathcal{A}v - v_t - v_a) = \mathcal{A}z - (z_t + z_a) - \frac{2m}{k} \lambda^{-1} z.$$

By the definition of inner product in $L^2(Q_{T,a^\dagger}^\Omega)$, we obtain

$$(3.6) \quad \begin{aligned} \|\lambda^{-\frac{m}{k}}(\mathcal{A}v - v_t - v_a)\|_{Q_{T,a^\dagger}^\Omega}^2 &= \|z_t + z_a\|_{Q_{T,a^\dagger}^\Omega}^2 - 2\langle z_t + z_a, \mathcal{A}z \rangle_{Q_{T,a^\dagger}^\Omega} \\ &+ \frac{4m}{k} \langle \lambda^{-1}z, z_t + z_a \rangle_{Q_{T,a^\dagger}^\Omega} + \left\| \mathcal{A}z - \frac{2m}{k} \lambda^{-1}z \right\|_{Q_{T,a^\dagger}^\Omega}^2. \end{aligned}$$

For the second term in right-hand side (RHS) of (3.6), using integration by parts

$$\begin{aligned} -2\langle z_t + z_a, \mathcal{A}z \rangle_{Q_{T,a^\dagger}^\Omega} &= 2 \int_0^T \int_0^{a^\dagger} \int_\Omega \sum_{1 \leq i, j \leq n} (\partial_t \partial_{x_i} z + \partial_a \partial_{x_i} z) d_{ij} \partial_{x_j} z dx dadt \\ &= 2 \int_0^T \int_0^{a^\dagger} \int_\Omega (\partial_t + \partial_a) \left(\sum_{1 \leq i, j \leq n} \partial_{x_i} z d_{ij} \partial_{x_j} z \right) dx dadt \\ &- 2 \int_0^T \int_0^{a^\dagger} \int_\Omega \sum_{1 \leq i, j \leq n} \partial_{x_i} z (\partial_t + \partial_a) (d_{ij} \partial_{x_j} z) dx dadt \\ &= -2 \int_0^T \int_0^{a^\dagger} \int_\Omega \sum_{1 \leq i, j \leq n} \partial_{x_i} z (\partial_t + \partial_a) (d_{ij}) \partial_{x_j} z dx dadt \\ &- 2 \int_0^T \int_0^{a^\dagger} \int_\Omega \sum_{1 \leq i, j \leq n} \partial_{x_i} z d_{ij} (\partial_t + \partial_a) \partial_{x_j} z dx dadt, \end{aligned}$$

where we have used that $z \in P_{T,a^\dagger}^\Omega$. Since $d_{ij} = d_{ji}$ (cf. (A₁)), from the first and fifth rows we obtain

$$\begin{aligned} &2 \int_0^T \int_0^{a^\dagger} \int_\Omega \sum_{1 \leq i, j \leq n} (\partial_t + \partial_a) (\partial_{x_i} z) d_{ij} \partial_{x_j} z dx dadt \\ &= - \int_0^T \int_0^{a^\dagger} \int_\Omega \sum_{1 \leq i, j \leq n} \partial_{x_i} z (\partial_t + \partial_a) (d_{ij}) \partial_{x_j} z dx dadt. \end{aligned}$$

Thus, we obtain that

$$(3.7) \quad -2\langle z_t + z_a, \mathcal{A}z \rangle_{Q_{T,a^\dagger}^\Omega} = - \int_0^T \int_0^{a^\dagger} \int_\Omega \sum_{1 \leq i, j \leq n} \partial_{x_i} z (\partial_t + \partial_a) (d_{ij}) \partial_{x_j} z dx dadt.$$

In the same manner, the third term in the RHS of (3.6) is

$$(3.8) \quad \begin{aligned} \frac{4m}{k} \langle \lambda^{-1}z, z_t + z_a \rangle_{Q_{T,a^\dagger}^\Omega} &= \frac{2m}{k} \int_0^T \int_0^{a^\dagger} \int_\Omega (\partial_t + \partial_a) (\lambda^{-1}z^2) dx dadt \\ &+ \frac{4m}{k} \int_0^T \int_0^{a^\dagger} \int_\Omega \lambda^{-2} z^2 dx dadt = \frac{4m}{k} \|\lambda^{-1}z\|_{Q_{T,a^\dagger}^\Omega}^2. \end{aligned}$$

Using (A₂), from (3.7) and (3.8), we obtain

$$\begin{aligned} \|\lambda^{-\frac{m}{k}}(\mathcal{A}v - v_t - v_a)\|_{Q_{T,a^\dagger}^\Omega}^2 &\geq \frac{4m}{k} \|\lambda^{-1}z\|_{Q_{T,a^\dagger}^\Omega}^2 \\ -\bar{M} \int_0^T \int_0^{a^\dagger} \int_\Omega \sum_{1 \leq i, j \leq n} |\partial_{x_i} z \partial_{x_j} z| dx dadt &\geq \frac{4m}{k} \|\lambda^{-1}z\|_{Q_{T,a^\dagger}^\Omega}^2 - \bar{M} \|\nabla z\|_{Q_{T,a^\dagger}^\Omega}^2. \end{aligned}$$

Substituting $z = \lambda^{-\frac{m}{k}} v$, we end the proof of (3.3).

Using (A₁), integration by parts and $|\lambda(t, a)| \leq T + a_{\dagger} + \eta$ we have

$$(3.9) \quad \begin{aligned} - \left\langle \lambda^{-\frac{2m}{k}} v, \mathcal{A}v - v_t - v_a \right\rangle_{Q_{T, a_{\dagger}}^{\Omega}} &\geq -\frac{2m}{k} (T + a_{\dagger} + \eta) \left\| \lambda^{-\frac{m}{k}-1} v \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 \\ &\quad + \underline{c} \left\| \lambda^{-\frac{m}{k}} \nabla v \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 \end{aligned}$$

It then follows from (3.3) that (3.9) can be estimated by

$$(3.10) \quad \begin{aligned} - \left\langle \lambda^{-\frac{2m}{k}} v, \mathcal{A}v - v_t - v_a \right\rangle_{Q_{T, a_{\dagger}}^{\Omega}} &\geq -\frac{(T + a_{\dagger} + \eta)}{2} \left(\left\| \lambda^{-\frac{m}{k}} (\mathcal{A}v - v_t - v_a) \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 \right. \\ &\quad \left. + \overline{M} \left\| \lambda^{-\frac{m}{k}} \nabla v \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 \right) + \underline{c} \left\| \lambda^{-\frac{m}{k}} \nabla v \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2. \end{aligned}$$

Using (3.3), we have

$$\begin{aligned} - \left\langle \lambda^{-\frac{2m}{k}} v, \mathcal{A}v - v_t - v_a \right\rangle_{Q_{T, a_{\dagger}}^{\Omega}} &\leq \frac{1}{2} \left\| \lambda^{-\frac{m}{k}-1} v \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 + \frac{1}{2} \left\| \lambda^{-\frac{m}{k}+1} (\mathcal{A}v - v_t - v_a) \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 \\ &\leq \frac{k}{8m} \left(\left\| \lambda^{-\frac{m}{k}} (\mathcal{A}v - v_t - v_a) \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 + \overline{M} \left\| \lambda^{-\frac{m}{k}} \nabla v \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 \right) \\ &\quad + \frac{(T + a_{\dagger} + \eta)^2}{2} \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}v - v_t - v_a) \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2. \end{aligned}$$

Combining this with (3.10) then reads as

$$\begin{aligned} &\underline{c} \left\| \lambda^{-\frac{m}{k}} \nabla v \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 \\ &\leq \left(\frac{k}{8m} + \frac{(T + a_{\dagger} + \eta)}{2} + \frac{(T + a_{\dagger} + \eta)^2}{2} \right) \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}v - v_t - v_a) \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 \\ &\quad + \left(\frac{(T + a_{\dagger} + \eta) \overline{M}}{2} + \frac{k \overline{M}}{8m} \right) \left\| \lambda^{-\frac{m}{k}} \nabla v \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2. \end{aligned}$$

Accordingly, if $\mu_0 \geq \mu \geq T + a_{\dagger} > 0$, $\eta_0 \geq \eta > 0$ and m_0 are such that

$$(3.11) \quad 2 \left(\frac{(\mu_0 + \eta_0) \overline{M}}{2} + \frac{k \overline{M}}{8m_0} \right) \leq \underline{c},$$

then for any $m \geq m_0$, we obtain

$$(3.12) \quad \left\| \lambda^{-\frac{m}{k}} \nabla v \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 \leq C_1 \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}v - v_t - v_a) \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2,$$

where $C_1 := \frac{2}{\underline{c}} \left(\frac{k}{8m} + \frac{(\mu+\eta)}{2} + \frac{(\mu+\eta)^2}{2} \right)$. Combining (3.12) and (3.3), we conclude that

$$\begin{aligned} &\left\| \lambda^{-\frac{m}{k}-1} v \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 + \frac{1}{2} \left\| \lambda^{-\frac{m}{k}} \nabla v \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 \\ &\leq \left[\frac{k}{4m} + C_1 \left(\frac{1}{2} + \frac{k \overline{M}}{4m} \right) \right] \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}v - v_t - v_a) \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2. \end{aligned}$$

Then denoting $C_2 = \frac{2}{\underline{c}} \left(\frac{k}{8m_0} + \frac{(\mu_0+\eta_0)}{2} + \frac{(\mu_0+\eta_0)^2}{2} \right)$ and choosing

$$(3.13) \quad K := \frac{k}{4m_0} + C_2 \left(\frac{1}{2} + \frac{k \overline{M}}{4m_0} \right),$$

we conclude that (3.4) holds for $m \geq m_0$. \square

Let us now choose m_0 sufficiently large, and μ_0 and η_0 sufficiently small such that K given by (3.13) satisfies

$$(3.14) \quad 0 < K \leq \frac{1}{4\alpha L_F^2}.$$

Let also $0 < \eta_1 \leq \min\{1, \eta_0\}$ and choose

$$(3.15) \quad \eta = \eta_1, \quad \mu'_0 = \eta^{\frac{m(\frac{1}{\alpha}-1)}{m(\frac{1}{\alpha}-1)+k}} 2^{\frac{k}{2m(\frac{1}{\alpha}-1)+2k}} - \eta > 0.$$

We consider here $T + a_\dagger \leq \mu'_0$ since the uniqueness result for the latter case $T + a_\dagger > \mu'_0$ is a direct consequence from the former case. In that case, the time and age intervals can be divided into many countable subsets whose lengths are not larger than μ'_0 , which brings us back to the former case.

Let $0 < t_1 < t_2 < T$ and $0 < a_1 < a_2 < a_\dagger$ and take $\kappa : \overline{Q_{T, a_\dagger}} \rightarrow \mathbb{R}$ such that it is twice continuously differentiable in $(t_1, t_2) \times (a_1, a_2)$ and

$$(3.16) \quad \kappa(t, a) = \begin{cases} 0 & \text{if } (t, a) \in Q_{T, a_\dagger} \setminus ((t_1, t_2) \times (a_1, a_2) \cup [t_2, T] \times [a_2, a_\dagger]), \\ 1 & \text{if } (t, a) \in [t_2, T] \times [a_2, a_\dagger]. \end{cases}$$

Let $v = \kappa w$, where w is the solution of (3.2), then notice that $v \in P_{T, a_\dagger}^\Omega$. Starting from the estimate (3.4), we have that for sufficiently large $m \geq m_0$

$$(3.17) \quad \begin{aligned} & K \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}v - v_t - v_a) \right\|_{Q_{(t_1, t_2) \times (a_1, a_2)}^\Omega}^2 + K \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}w - w_t - w_a) \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2 \\ & \geq \left\| \lambda^{-\frac{m}{k}-1} w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2 + \frac{1}{2} \left\| \lambda^{-\frac{m}{k}} \nabla v \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2. \end{aligned}$$

From (3.2) and (A₃), we get $\lambda^{-\frac{2m}{k}} (\mathcal{A}w - w_t - w_a)^2 \leq \lambda^{-\frac{2m}{k}} L_F^2 |w|^{2\alpha}$, then applying Lemma 2.1 with $X = |w|^2 > 0$, $\alpha_0 = \alpha$ and

$$(3.18) \quad \gamma = [(t_2 - t_1)(a_2 - a_1)]^{-\frac{1}{\alpha}} \lambda^{\frac{2m}{k\alpha}} \left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^{\frac{2}{\alpha}} > 0,$$

we have

$$\begin{aligned} \lambda^{-\frac{2m}{k}} (\mathcal{A}w - w_t - w_a)^2 & \leq \alpha L_F^2 \frac{\lambda^{-\frac{2m}{k\alpha}} \left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^{\frac{2}{\alpha}(\alpha-1)}}{[(t_2 - t_1)(a_2 - a_1)]^{1-\frac{1}{\alpha}}} w^2 \\ & \quad + (1 - \alpha) L_F^2 \frac{\left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^2}{(t_2 - t_1)(a_2 - a_1)}. \end{aligned}$$

Integrating both sides over Ω , we have

$$(3.19) \quad \begin{aligned} \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}w - w_t - w_a) \right\|_\Omega^2 & \leq \alpha L_F^2 \frac{\lambda^{-\frac{2m}{k\alpha} + \frac{2m}{k}} \left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^{\frac{2}{\alpha}(\alpha-1)}}{[(t_2 - t_1)(a_2 - a_1)]^{1-\frac{1}{\alpha}}} \lambda^{-\frac{2m}{k}} \|w\|_\Omega^2 \\ & \quad + (1 - \alpha) |\Omega| L_F^2 \frac{\left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^2}{(t_2 - t_1)(a_2 - a_1)}. \end{aligned}$$

Notice that for $\alpha \in (0, 1]$, it holds $\lambda^{-\frac{2m}{k\alpha} + \frac{2m}{k}} = \lambda^{\frac{2m}{k}(1-\frac{1}{\alpha})} \leq \eta^{\frac{2m}{k}(1-\frac{1}{\alpha})}$. Using this bound and integrating both sides of (3.19) over $(t_2, T) \times (a_2, a_\dagger)$, we obtain

$$\begin{aligned}
& \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}w - w_t - w_a) \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2 \\
& \leq \alpha L_F^2 \frac{\left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^{2(1-\frac{1}{\alpha})}}{(t_2 - t_1)^{1-\frac{1}{\alpha}} (a_2 - a_1)^{1-\frac{1}{\alpha}}} \left\| \lambda^{-\frac{m}{k}} w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2 \eta^{\frac{2m}{k}(1-\frac{1}{\alpha})} \\
(3.20) \quad & + (1 - \alpha) |\Omega| \frac{(T - t_2)(a_\dagger - a_2)}{(t_2 - t_1)(a_2 - a_1)} L_F^2 \left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^2.
\end{aligned}$$

Using that $|\lambda| \leq \mu'_0 + \eta$, we have

$$(3.21) \quad \left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^{2(1-\frac{1}{\alpha})} \leq (t_2 - t_1)^{1-\frac{1}{\alpha}} (a_2 - a_1)^{1-\frac{1}{\alpha}} (\mu'_0 + \eta)^{\frac{2m}{k}(\frac{1}{\alpha}-1)},$$

$$(3.22) \quad \left\| \lambda^{-\frac{m}{k}} w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2 \leq (\mu'_0 + \eta)^2 \left\| \lambda^{-\frac{m}{k}-1} w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2.$$

Plugging (3.20)-(3.22) into (3.17) and by the choice (3.15), we obtain

$$\begin{aligned}
(3.23) \quad & K \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}v - v_t - v_a) \right\|_{Q_{(t_1, t_2) \times (a_1, a_2)}^\Omega}^2 \\
& + K |\Omega| \frac{(T - t_2)(a_\dagger - a_2)}{(t_2 - t_1)(a_2 - a_1)} L_F^2 \left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^2 \geq \frac{1}{2} \left\| \lambda^{-\frac{m}{k}-1} w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2.
\end{aligned}$$

For any $t_2 < t_3 < T$ and $a_2 < a_3 < a_\dagger$, (3.23) can be further estimated by

$$\begin{aligned}
(3.24) \quad & K (T + a_\dagger + \eta - t_2 - a_2)^{-\frac{2m}{k}} \left(\left\| \mathcal{A}v - v_t - v_a \right\|_{Q_{(t_1, t_2) \times (a_1, a_2)}^\Omega}^2 + T a_\dagger |\Omega| L_F^2 \right) \\
& \geq \frac{(T + a_\dagger + \eta - t_3 - a_3)^{-\frac{2m}{k}-2}}{2} \left\| w \right\|_{Q_{(t_3, T) \times (a_3, a_\dagger)}^\Omega}^2.
\end{aligned}$$

Observing that

$$(3.25) \quad \left(\frac{T + a_\dagger + \eta - t_2 - a_2}{T + a_\dagger + \eta - t_3 - a_3} \right)^{-\frac{2m}{k}} \rightarrow 0 \quad \text{as } m \rightarrow \infty,$$

we obtain from (3.24) that $w \equiv 0$ for $(t, a, x) \in Q_{(t_3, T) \times (a_3, a_\dagger)}^\Omega$ and thus for $(t, a, x) \in Q_{T, a_\dagger}^\Omega$ since $0 < t_1 < t_2 < t_3$ and $0 < a_1 < a_2 < a_3$ can be taken arbitrarily small.

Hence, we state the following uniqueness theorem.

Theorem 3.2. *Assume (A₁)-(A₃) hold. Then, the problem (1.4) with the Dirichlet boundary condition (1.5) admits no more than one solution in W_{T, a_\dagger}^Ω .*

Remark 3.3. The presence of $k > 0$ in the context of Lemma 3.1 implies that m can be taken as a real positive number. As a corollary, the uniqueness result holds when the source term F is globally Lipschitz, i.e. $\alpha = 1$. When F is locally Lipschitz, it can also be reduced to the global case if we set

$$\widehat{W}_{T, a_\dagger}^\Omega = C(\overline{Q_{T, a_\dagger}}; H_0^1(\Omega) \cap L^\infty(\Omega)) \cap L^2(Q_{T, a_\dagger}; H^2(\Omega)) \cap C^1(Q_{T, a_\dagger}; H_0^1(\Omega)).$$

This space implies that the population density is essentially bounded in space and uniformly bounded in time and age. This further gives us an idea to obtain the uniqueness of solution when $\alpha > 1$. In fact, if we assign this boundedness to a constant $r > 0$ and apply the elementary inequality $|X|^\alpha \leq \alpha r^{\alpha-1} |X|$ provided

that $|X| \leq r$ and $\alpha > 1$, then we get back the globally Lipschitz case.

Concerning the existence of the function κ in (3.16), we can rely on the application of partitions of unity (cf. [5, Proposition 2.25]). It says that if M is a smooth manifold with or without boundary, then for any closed subset $A \subseteq M$ and any open subset U containing A , there exists a smooth bump function for A supported in U . Accordingly, the existence of κ is guaranteed by taking,

$$A = [t_2, T] \times [a_2, a_\dagger], \quad M = \overline{Q_{T, a_\dagger}} \quad \text{and} \quad U = \{(t_1, t_2) \times (a_1, a_2)\} \cup A.$$

3.2. Robin-type boundary condition. Similar to the previous subsection, to prove uniqueness we consider u_1 and u_2 as two solutions (in some appropriate spaces) of (1.4) and (1.6) and then denote $w = u_1 - u_2$, which satisfies

$$(3.26) \quad \begin{cases} w_t + w_a - \mathcal{A}w = F(t, a, x; u_1) - F(t, a, x; u_2) & \text{in } Q_{T, a_\dagger}^\Omega, \\ -\text{d}(t, a, x) \nabla w(t, a, x) \cdot \mathbf{n} = S(u_1) - S(u_2) & \text{on } Q_{T, a_\dagger}^{\partial\Omega}, \\ w(T, a, x) = 0 & \text{in } (0, a_\dagger) \times \Omega, \\ w(t, a_\dagger, x) = 0 & \text{in } (0, T) \times \Omega. \end{cases}$$

We set the following function spaces:

$$(3.27) \quad \widetilde{W}_{T, a_\dagger}^\Omega := C(\overline{Q_{T, a_\dagger}}; H^1(\Omega)) \cap L^2(Q_{T, a_\dagger}; H^2(\Omega)) \cap C^1(Q_{T, a_\dagger}; H^1(\Omega)),$$

$$(3.28) \quad \widetilde{P}_{T, a_\dagger}^\Omega := \left\{ u \in \widetilde{W}_{T, a_\dagger}^\Omega : u|_{t \in \{0, T\}} = 0, u|_{a \in \{0, a_\dagger\}} = 0 \right\}.$$

The space $\widetilde{P}_{T, a_\dagger}^\Omega$ does not now contain any boundary information on $\partial\Omega$ as in P_{T, a_\dagger}^Ω due to the Robin-type boundary condition (1.6) instead of the Dirichlet boundary condition (1.5). Thus, Lemma 3.1 cannot be applied with any function in $\widetilde{P}_{T, a_\dagger}^\Omega$. However, we are able to formulate the following lemma with $w \in \widetilde{P}_{T, a_\dagger}^\Omega$ directly as a solution of (3.26). Here, we also set $\lambda(t, a) := t - T + a - a_\dagger - \eta$ for $\eta > 0$.

Lemma 3.4. *Let $\beta \in (0, 1)$ and assume (A_1) , (A_2) and (A_4) - (A_6) hold. Let $w \in \widetilde{P}_{T, a_\dagger}^\Omega$ be a solution of the problem (3.26). Then, for any $m \in \mathbb{N}^*$, $k \in \mathbb{R}_+^*$ and $0 < t_1 < t_2 < T$, $0 < a_1 < a_2 < a_\dagger$, we have*

$$(3.29) \quad \begin{aligned} & \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}w - w_t - w_a) \right\|_{Q_{T, a_\dagger}^\Omega}^2 \\ & \geq \frac{4m}{k} \left\| \lambda^{-\frac{m}{k} - 1} w \right\|_{Q_{T, a_\dagger}^\Omega}^2 - \overline{M} \left\| \lambda^{-\frac{m}{k}} \nabla w \right\|_{Q_{T, a_\dagger}^\Omega}^2 - 2\overline{m} \left\| \lambda^{-\frac{m}{k}} w \right\|_{Q_{T, a_\dagger}^{\partial\Omega}}^2 \\ & - \frac{\eta^2}{4C_0} \left\| \lambda^{-\frac{m}{k} - 1} w \right\|_{Q_{T, a_\dagger}^{\partial\Omega}}^2 - 16C_0 \mathbf{A}, \end{aligned}$$

where

$$(3.30) \quad \begin{aligned} \mathbf{A} & = \beta L_S^2 \frac{\eta^{\frac{2m}{k}(1-\frac{1}{\beta})+2} \left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^{\frac{2}{\beta}(\beta-1)}}{[(t_2 - t_1)(a_2 - a_1)]^{1-\frac{1}{\beta}}} \left\| \lambda^{-\frac{m}{k}} w \right\|_{Q_{T, a_\dagger}^{\partial\Omega}}^2 \\ & + \left(\frac{m}{k\eta^{\beta+1}} \right)^{\frac{2}{1-\beta}} (1-\beta) |\partial\Omega| L_S^2 \frac{T a_\dagger \left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^2}{(t_2 - t_1)(a_2 - a_1)}. \end{aligned}$$

Also, if $0 < T + a_{\dagger} \leq \mu$ for sufficiently small $\mu > 0$, there exists $K > 0$ such that

(3.31)

$$K \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}w - w_t - w_a) \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 + K \mathbf{A} \geq \left\| \lambda^{-\frac{m}{k}-1} w \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 + \frac{1}{2} \left\| \lambda^{-\frac{m}{k}} \nabla w \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2,$$

for sufficiently large m .

Proof. We can adapt the proof of Lemma 3.1 to prove the estimates (3.29) and (3.31). Omitting some calculus, let $z = \lambda^{-\frac{m}{k}} w \in \widetilde{P}_{T, a_{\dagger}}^{\Omega}$, then (as in (3.6))

$$(3.32) \quad \begin{aligned} \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}w - w_t - w_a) \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 &= \|z_t + z_a\|_{Q_{T, a_{\dagger}}^{\Omega}}^2 - 2 \langle z_t + z_a, \mathcal{A}z \rangle_{Q_{T, a_{\dagger}}^{\Omega}} \\ &\quad + \frac{4m}{k} \langle \lambda^{-1} z, z_t + z_a \rangle_{Q_{T, a_{\dagger}}^{\Omega}} + \left\| \mathcal{A}z - \frac{2m}{k} \lambda^{-1} z \right\|_{Q_{T, a_{\dagger}}^{\Omega}}^2. \end{aligned}$$

Using the Robin boundary condition from (3.26), we obtain

$$(3.33) \quad \begin{aligned} -2 \langle z_t + z_a, \mathcal{A}z \rangle_{L^2(Q_{T, a_{\dagger}}^{\Omega})} &= \underbrace{-2 \int_0^T \int_0^{a_{\dagger}} \int_{\partial\Omega} (z_t + z_a) d\nabla z \cdot n d\sigma_x dadt}_{:=\mathcal{I}_1} \\ &\quad + \underbrace{2 \int_0^T \int_0^{a_{\dagger}} \int_{\Omega} \sum_{1 \leq i, j \leq n} (\partial_t \partial_{x_i} z + \partial_a \partial_{x_i} z) d_{ij} \partial_{x_j} z dx dadt}_{:=\mathcal{I}_2}. \end{aligned}$$

As in the derivation of (3.7) and using (A₂) we obtain

$$(3.34) \quad \begin{aligned} \mathcal{I}_2 &= - \int_0^T \int_0^{a_{\dagger}} \int_{\Omega} \sum_{1 \leq i, j \leq n} \partial_{x_i} z (\partial_t + \partial_a) (d_{ij}) \partial_{x_j} z dx dadt \\ &\geq -\overline{M} \|\nabla z\|_{L^2(Q_{T, a_{\dagger}}^{\Omega})}^2. \end{aligned}$$

To estimate \mathcal{I}_1 , recall the Robin boundary condition from (3.26) to get

$$(3.35) \quad \begin{aligned} \mathcal{I}_1 &= 2 \int_0^T \int_0^{a_{\dagger}} \int_{\partial\Omega} (z_t + z_a) \lambda^{-\frac{m}{k}} (S(u_1) - S(u_2)) d\sigma_x dadt \\ &= 2 \underbrace{\int_0^T \int_0^{a_{\dagger}} \int_{\partial\Omega} (\partial_t + \partial_a) [\lambda^{-\frac{m}{k}} z (S(u_1) - S(u_2))] d\sigma_x dadt}_{:=\mathcal{I}_3} \\ &\quad - 2 \underbrace{\int_0^T \int_0^{a_{\dagger}} \int_{\partial\Omega} \lambda^{-\frac{m}{k}} z (\partial_t + \partial_a) (S(u_1) - S(u_2)) d\sigma_x dadt}_{:=\mathcal{I}_4} \\ &\quad + \underbrace{\frac{4m}{k} \int_0^T \int_0^{a_{\dagger}} \int_{\partial\Omega} \lambda^{-\frac{m}{k}-1} z (S(u_1) - S(u_2)) d\sigma_x dadt}_{:=\mathcal{I}_5}. \end{aligned}$$

Notice that due to the zero conditions in the definition (3.28) of $\tilde{P}_{T,a_\dagger}^\Omega$, the first term \mathcal{I}_3 of (3.35) vanishes. Using (A₄), we estimate \mathcal{I}_4 by

$$(3.36) \quad \begin{aligned} \mathcal{I}_4 &= -2 \int_0^T \int_0^{a_\dagger} \int_{\partial\Omega} \lambda^{-\frac{2m}{k}} (u_1 - u_2) (\partial_t + \partial_a) (S(u_1) - S(u_2)) d\sigma_x dadt \\ &\geq -2\bar{m} \|\lambda^{-\frac{m}{k}} w\|_{Q_{T,a_\dagger}^{\partial\Omega}}^2. \end{aligned}$$

By the back-substitution $z = \lambda^{-\frac{m}{k}} w$, the term \mathcal{I}_5 can be estimated by

$$(3.37) \quad \mathcal{I}_5 \geq -\frac{2m}{k} \left(\frac{k\eta^2}{8mC_0} \|\lambda^{-\frac{m}{k}-1} w\|_{Q_{T,a_\dagger}^{\partial\Omega}}^2 + \frac{8mC_0}{k\eta^2} \|\lambda^{-\frac{m}{k}} (S(u_1) - S(u_2))\|_{Q_{T,a_\dagger}^{\partial\Omega}}^2 \right).$$

Apply (A₆) and the inequality (2.1) for $X = |w|^2 \geq 0$, $\alpha_0 = \beta \in (0, 1)$ and

$$\gamma = [(t_2 - t_1)(a_2 - a_1)]^{-\frac{1}{\beta}} \left(\frac{\eta^2 k}{m} \right)^{\frac{2}{\beta-1}} \lambda^{\frac{2m}{k\beta}} \|\lambda^{-\frac{m}{k}}\|_{(t_1, t_2) \times (a_1, a_2)}^{\frac{2}{\beta}} > 0,$$

for $0 < t_1 < t_2 < T$, $0 < a_1 < a_2 < a_\dagger$, to get (using that $|\lambda| > \eta$), as in (3.19),

$$(3.38) \quad \frac{m^2}{k^2 \eta^2} \|\lambda^{-\frac{m}{k}} (S(u_1) - S(u_2))\|_{Q_{T,a_\dagger}^{\partial\Omega}}^2 \leq \mathbf{A}.$$

As in (3.8) since $z \in \tilde{P}_{T,a_\dagger}^\Omega$, we observe that

$$(3.39) \quad \frac{4m}{k} \langle \lambda^{-1} z, z_t + z_a \rangle_{Q_{T,a_\dagger}^\Omega} = \frac{4m}{k} \|\lambda^{-1} z\|_{Q_{T,a_\dagger}^\Omega}^2.$$

We complete the proof of (3.29) by grouping together (3.32), (3.34), (3.36)-(3.39). It remains to prove the estimate (3.31). Using Green's formula we have

$$(3.40) \quad \begin{aligned} -\langle \lambda^{-\frac{2m}{k}} w, \mathcal{A}w - w_t - w_a \rangle_{Q_{T,a_\dagger}^\Omega} &= \underbrace{\int_0^T \int_0^{a_\dagger} \int_{\partial\Omega} \lambda^{-\frac{2m}{k}} w (S(u_1) - S(u_2)) d\sigma_x dadt}_{:=\mathcal{I}_6} \\ &\quad + \underbrace{\int_0^T \int_0^{a_\dagger} \int_{\Omega} \lambda^{-\frac{2m}{k}} (d\nabla w) \cdot \nabla w dx dadt}_{:=\mathcal{I}_7} \\ &\quad + \underbrace{\int_0^T \int_0^{a_\dagger} \int_{\Omega} \lambda^{-\frac{2m}{k}} w (w_t + w_a) dx dadt}_{:=\mathcal{I}_8}. \end{aligned}$$

Since $|\lambda(t, a)| \leq T + a_\dagger + \eta$ for all $(t, a) \in \overline{Q_{T,a_\dagger}}$, \mathcal{I}_8 can be estimated using integration by parts, while for \mathcal{I}_6 and \mathcal{I}_7 we use (A₁) and (A₅) to obtain

$$(3.41) \quad \begin{aligned} -\langle \lambda^{-\frac{2m}{k}} w, \mathcal{A}w - w_t - w_a \rangle_{Q_{T,a_\dagger}^\Omega} &\geq \underline{c} \|\lambda^{-\frac{m}{k}} \nabla w\|_{Q_{T,a_\dagger}^\Omega}^2 \\ &\quad - \frac{2m}{k} (T + a_\dagger + \eta) \|\lambda^{-\frac{m}{k}-1} w\|_{Q_{T,a_\dagger}^\Omega}^2. \end{aligned}$$

Using Young's inequality

$$\begin{aligned} & \frac{1}{2} \|\lambda^{-\frac{m}{k}-1} w\|_{Q_{T,a_\dagger}^\Omega}^2 + \frac{1}{2} \|\lambda^{-\frac{m}{k}+1} (\mathcal{A}w - w_t - w_a)\|_{Q_{T,a_\dagger}^\Omega}^2 \\ & \geq - \left\langle \lambda^{-\frac{2m}{k}} w, \mathcal{A}w - w_t - w_a \right\rangle_{Q_{T,a_\dagger}^\Omega}, \end{aligned}$$

then (3.41) yields (using also that $|\lambda| \leq T + a_\dagger + \eta$)

$$(3.42) \quad \begin{aligned} & K_2 \|\lambda^{-\frac{m}{k}-1} w\|_{Q_{T,a_\dagger}^\Omega}^2 + \frac{(T + a_\dagger + \eta)^2}{2} \|\lambda^{-\frac{m}{k}} (\mathcal{A}w - w_t - w_a)\|_{Q_{T,a_\dagger}^\Omega}^2 \\ & \geq \underline{c} \|\lambda^{-\frac{m}{k}} \nabla w\|_{Q_{T,a_\dagger}^\Omega}^2 \end{aligned}$$

where $K_2 := \frac{1}{2} + \frac{2m}{k} (T + a_\dagger + \eta)$. Now using (2.2) in the third and fourth terms of the RHS of (3.29), we obtain (using also that $|\lambda| > \eta$)

$$(3.43) \quad \begin{aligned} & \|\lambda^{-\frac{m}{k}} (\mathcal{A}w - w_t - w_a)\|_{Q_{T,a_\dagger}^\Omega}^2 \\ & \geq K_1 \|\lambda^{-\frac{m}{k}-1} w\|_{Q_{T,a_\dagger}^\Omega}^2 - \left(\overline{M} + 2\overline{m}C_0 + \frac{1}{4} \right) \|\lambda^{-\frac{m}{k}} \nabla w\|_{Q_{T,a_\dagger}^\Omega}^2 - 16C_0 \mathbf{A}, \end{aligned}$$

where $K_1 := \frac{4m}{k} - 2\overline{m}C_0 (T + a_\dagger + \eta)^2 - \frac{\eta^2}{4}$. Multiplying (3.43) by $K_2 K_1^{-1}$ and applying (3.42), we obtain

$$(3.44) \quad \left[\underline{c} - K_2 K_1^{-1} \left(\overline{M} + 2C_0 \overline{m} + \frac{1}{4} \right) \right] \|\lambda^{-\frac{m}{k}} \nabla w\|_{Q_{T,a_\dagger}^\Omega}^2 \leq \mathcal{I}_9 + \mathcal{I}_{10},$$

which in line with the estimate (3.12). In (3.44), we have denoted

$$\begin{aligned} \mathcal{I}_9 & := \left[K_2 K_1^{-1} + \frac{(T + a_\dagger + \eta)^2}{2} \right] \|\lambda^{-\frac{m}{k}} (\mathcal{A}w - w_t - w_a)\|_{Q_{T,a_\dagger}^\Omega}^2, \\ \mathcal{I}_{10} & := 16K_2 K_1^{-1} C_0 \mathbf{A}. \end{aligned}$$

Since $K_2 K_1^{-1} \rightarrow \frac{1}{2} (T + a_\dagger + \eta)$ as $m \rightarrow \infty$, if $0 < T + a_\dagger \leq \mu$ for sufficiently small $\mu > 0$, then can choose $\mu_0 \geq \mu \geq T + a_\dagger > 0$, $\eta_0 \geq \eta > 0$ and $m_0 \leq m$ such that $K_1 > 0$ and $2K_2 K_1^{-1} (\overline{M} + 2C_0 \overline{m} + \frac{1}{4}) \leq \underline{c}$. Then, for any $m \geq m_0$ denoting

$$C_3 := \frac{2}{\underline{c}} \max \left\{ K_2 K_1^{-1} + \frac{(\mu + \eta)^2}{2}, 16K_2 K_1^{-1} C_0 \right\},$$

from (3.43) and (3.44) we obtain

$$(3.45) \quad \begin{aligned} & \|\lambda^{-\frac{m}{k}-1} w\|_{Q_{T,a_\dagger}^\Omega}^2 + \left[1 - K_1^{-1} \left(\overline{M} + 2C_0 \overline{m} + \frac{1}{4} \right) \right] \|\lambda^{-\frac{m}{k}} \nabla w\|_{Q_{T,a_\dagger}^\Omega}^2 \\ & \leq (K_1^{-1} + C_3) \left(\|\lambda^{-\frac{m}{k}} (\mathcal{A}w - w_t - w_a)\|_{Q_{T,a_\dagger}^\Omega}^2 + \mathbf{A} \right) \end{aligned}$$

We can now take m_0 sufficiently large, and μ_0 and η_0 sufficiently small such that $K_1^{-1} (\overline{M} + 2C_0 \overline{m} + \frac{1}{4}) \leq \frac{1}{2}$ for any $m \geq m_0$. Then, we conclude that (3.31) holds

for $m \geq m_0$ by choosing $K := K_3^{-1} + C_4$, where

$$K_3 := \frac{4m_0}{k} - \frac{\eta^2}{4} - 2\bar{m}(\mu_0 + \eta_0)^2 C_0, \quad K_4 := \frac{1}{2} + \frac{2m_0}{k}(\mu_0 + \eta_0),$$

$$C_4 := \frac{2}{\underline{c}} \max \left\{ K_4 K_3^{-1} + \frac{(\mu_0 + \eta_0)^2}{2}, 16K_4 K_3^{-1} C_0 \right\},$$

using that $K_4 K_3^{-1} \geq K_2 K_1^{-1}$ since the function $K_2 K_1^{-1}$ is monotonically decreasing, as a function of m . \square

Remark that in deriving (3.29) and (3.31), we have never used that w satisfies the first equation in (3.26). Let us now define

$$\bar{\kappa}(t, a) = \begin{cases} 0 & \text{if } (t, a) \in Q_{T, a_\dagger} \setminus ((t_1, t_2) \times (a_1, a_2)) \cup [t_2, T] \times [a_2, a_\dagger], \\ 1 & \text{if } (t, a) \in ((t_1, t_2) \times (a_1, a_2)) \cup [t_2, T] \times [a_2, a_\dagger], \end{cases}$$

and set $v := \bar{\kappa}w \in \tilde{P}_{T, a_\dagger}^\Omega$. Then, from the second equation of (3.26) we have

$$\begin{aligned} & -d(t, a, x) \nabla v(t, a, x) \cdot \mathbf{n} \\ &= \begin{cases} S(u_1) - S(u_2) & \text{on } ((t_1, t_2) \times (a_1, a_2) \cup [t_2, T] \times [a_2, a_\dagger]) \times \partial\Omega, \\ 0 & \text{on } (Q_{T, a_\dagger} \setminus ((t_1, t_2) \times (a_1, a_2)) \cup [t_2, T] \times [a_2, a_\dagger]) \times \partial\Omega. \end{cases} \end{aligned}$$

Then, it can be remarked that the inequality (3.31) in Lemma 3.4 holds not only for w but also for $v = \bar{\kappa}w$ since in (3.40) we only need to use that $\mathcal{I}_6 \geq 0$, with the rest of the argument remaining the same. Then, from (3.31) applied to v we have (since $v = w$ in $((t_1, t_2) \times (a_1, a_2) \cup [t_2, T] \times [a_2, a_\dagger]) \times \Omega$ and $v = 0$ otherwise)

(3.46)

$$\begin{aligned} & K \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}w - w_t - w_a) \right\|_{Q_{(t_1, t_2) \times (a_1, a_2)}^\Omega}^2 + K \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}w - w_t - w_a) \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2 \\ &+ K \mathbf{A}_1 + K \mathbf{A}_2 \geq \left\| \lambda^{-\frac{m}{k}-1} w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2 + \frac{1}{2} \left\| \lambda^{-\frac{m}{k}} \nabla w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2, \end{aligned}$$

where

$$\begin{aligned} \mathbf{A}_1 &= \beta L_S^2 \frac{\eta^{\frac{2m}{k}(1-\frac{1}{\beta})+2} \left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^{\frac{2}{\beta}(\beta-1)}}{[(t_2 - t_1)(a_2 - a_1)]^{1-\frac{1}{\beta}}} \left\| \lambda^{-\frac{m}{k}} w \right\|_{Q_{(t_1, t_2) \times (a_1, a_2)}^{\partial\Omega}}^2 \\ &+ \left(\frac{m}{k\eta^{\beta+1}} \right)^{\frac{2}{1-\beta}} (1-\beta) |\partial\Omega| L_S^2 \frac{T a_\dagger \left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^2}{(t_2 - t_1)(a_2 - a_1)}, \\ \mathbf{A}_2 &= \beta L_S^2 \frac{\eta^{\frac{2m}{k}(1-\frac{1}{\beta})+2} \left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^{\frac{2}{\beta}(\beta-1)}}{[(t_2 - t_1)(a_2 - a_1)]^{1-\frac{1}{\beta}}} \left\| \lambda^{-\frac{m}{k}} w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^{\partial\Omega}}^2. \end{aligned}$$

Since $|\lambda| \leq T + a_\dagger + \eta \leq \mu'_0 + \eta$, we have that

$$(3.47) \quad \begin{aligned} \mathbf{A}_1 &\leq \beta L_S^2 (\mu'_0 + \eta)^{\frac{2m}{k}(\frac{1}{\beta}-1)+2} \eta^{\frac{2m}{k}(1-\frac{1}{\beta})+2} \left\| \lambda^{-\frac{m}{k}-1} w \right\|_{Q_{(t_1, t_2) \times (a_1, a_2)}^{\partial\Omega}}^2 \\ &+ \left(\frac{m}{k\eta^{\beta+1}} \right)^{\frac{2}{1-\beta}} (1-\beta) |\partial\Omega| L_S^2 \frac{T a_\dagger \left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^2}{(t_2 - t_1)(a_2 - a_1)}. \end{aligned}$$

Also, as in (3.21), with α replaced by β , using (2.2) and $|\lambda| \leq \mu'_0 + \eta$, we have

$$(3.48) \quad \mathbf{A}_2 \leq \beta L_S^2 (\mu'_0 + \eta)^{\frac{2m}{k}(\frac{1}{\beta}-1)+2} \eta^{\frac{2m}{k}(1-\frac{1}{\beta})+2} C_0 \left\| \lambda^{-\frac{m}{k}-1} w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2 \\ + \beta L_S^2 (\mu'_0 + \eta)^{\frac{2m}{k}(\frac{1}{\beta}-1)} \eta^{\frac{2m}{k}(1-\frac{1}{\beta})+2} C_0 \left\| \lambda^{-\frac{m}{k}} \nabla w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2.$$

From (3.20)-(3.22), we have that

$$(3.49) \quad \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}w - w_t - w_a) \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2 \\ \leq \alpha L_F^2 \frac{\eta^{\frac{2m}{k}(1-\frac{1}{\alpha})} \left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^{\frac{2}{\alpha}(\alpha-1)}}{[(t_2 - t_1)(a_2 - a_1)]^{1-\frac{1}{\alpha}}} \left\| \lambda^{-\frac{m}{k}} w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2 \\ + (1 - \alpha) |\Omega| L_F^2 \frac{Ta_\dagger \left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^2}{(t_2 - t_1)(a_2 - a_1)} \\ \leq \alpha L_F^2 (\mu'_0 + \eta)^{\frac{2m}{k}(\frac{1}{\alpha}-1)+2} \eta^{\frac{2m}{k}(1-\frac{1}{\alpha})} \left\| \lambda^{-\frac{m}{k}-1} w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2 \\ + (1 - \alpha) |\Omega| L_F^2 \frac{Ta_\dagger \left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^2}{(t_2 - t_1)(a_2 - a_1)}.$$

Combining (3.47)-(3.48), for sufficiently large $m \geq m_0$ it follows from (3.46) (noticing also $\eta^2 \leq (\mu'_0 + \eta)^2$) that

$$(3.50) \quad K \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}w - w_t - w_a) \right\|_{Q_{(t_1, t_2) \times (a_1, a_2)}^\Omega}^2 \\ + K \beta L_S^2 (\mu'_0 + \eta)^{\frac{2m}{k}(\frac{1}{\beta}-1)+2} \eta^{\frac{2m}{k}(1-\frac{1}{\beta})} \left(\left\| \lambda^{-\frac{m}{k}-1} w \right\|_{Q_{(t_1, t_2) \times (a_1, a_2)}^{\beta\Omega}}^2 \right. \\ \left. + C_0 \left\| \lambda^{-\frac{m}{k}} \nabla w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2 \right) \\ + K \left[\alpha L_F^2 (\mu'_0 + \eta)^{\frac{2m}{k}(\frac{1}{\alpha}-1)+2} \eta^{\frac{2m}{k}(1-\frac{1}{\alpha})} \right. \\ \left. + C_0 \beta L_S^2 (\mu'_0 + \eta)^{\frac{2m}{k}(\frac{1}{\beta}-1)+2} \eta^{\frac{2m}{k}(1-\frac{1}{\beta})} \right] \left\| \lambda^{-\frac{m}{k}-1} w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2 \\ + K \frac{Ta_\dagger}{(t_2 - t_1)(a_2 - a_1)} \left[(1 - \alpha) |\Omega| L_F^2 \right. \\ \left. + (1 - \beta) |\partial\Omega| L_S^2 \left(\frac{m}{k\eta^{\beta+1}} \right)^{\frac{2}{1-\beta}} \right] \left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^2 \\ \geq \left\| \lambda^{-\frac{m}{k}-1} w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2 + \frac{1}{2} \left\| \lambda^{-\frac{m}{k}} \nabla w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2.$$

Let $0 < \eta_1 \leq \min\{1, \eta_0\}$, take $\eta = \eta_1$ and

$$(3.51) \quad \mu'_0 = \begin{cases} \eta^{\frac{m(\frac{1}{\beta}-1)}{m(\frac{1}{\beta}-1)+k}} 2^{\frac{k}{2m(\frac{1}{\beta}-1)+2k}} - \eta & \text{if } \alpha \geq \beta, \\ \eta^{\frac{m(\frac{1}{\alpha}-1)}{m(\frac{1}{\alpha}-1)+k}} 2^{\frac{k}{2m(\frac{1}{\alpha}-1)+2k}} - \eta & \text{if } \alpha < \beta. \end{cases}$$

Also, for sufficiently large m_0 , and μ_0, η_0 sufficiently small, choose

$$(3.52) \quad 0 < K \leq \frac{1}{8} \min \left\{ \frac{1}{\alpha L_F^2}, \frac{1}{C_0 \beta L_S^2}, \frac{1}{\beta L_S^2} \right\}.$$

With the choice (3.51), we have that $(\mu'_0 + \eta)^{\frac{2m}{k}(\frac{1}{\beta}-1)+2} \eta^{\frac{2m}{k}(1-\frac{1}{\beta})} = 2$. Further, with the choice (3.52) we have that $2K(\alpha L_F^2 + C_0 \beta L_S^2) \leq \frac{1}{2}$ and then (3.50) yields

$$\begin{aligned} & K \left\| \lambda^{-\frac{m}{k}} (\mathcal{A}w - w_t - w_a) \right\|_{Q_{(t_1, t_2) \times (a_1, a_2)}^\Omega}^2 + \frac{1}{4} \left\| \lambda^{-\frac{m}{k}-1} w \right\|_{Q_{(t_1, t_2) \times (a_1, a_2)}^{\partial\Omega}}^2 \\ & + \frac{KTa_\dagger}{(t_2 - t_1)(a_2 - a_1)} \left[|\Omega| L_F^2 + |\partial\Omega| L_S^2 \left(\frac{m}{k\eta^{\beta+1}} \right)^{\frac{2}{1-\beta}} \right] \left\| \lambda^{-\frac{m}{k}} \right\|_{(t_1, t_2) \times (a_1, a_2)}^2 \\ & \geq \frac{1}{2} \left\| \lambda^{-\frac{m}{k}-1} w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2 + \frac{1}{4} \left\| \lambda^{-\frac{m}{k}} \nabla w \right\|_{Q_{(t_2, T) \times (a_2, a_\dagger)}^\Omega}^2. \end{aligned}$$

Observe that to complete the uniqueness result, we only need to mimic the way (3.24) was obtained from (3.23), and the limit

$$\left(\frac{m}{k} \right)^{\frac{2}{1-\beta}} \left(\frac{T + a_\dagger + \eta - t_2 - a_2}{T + a_\dagger + \eta - t_3 - a_3} \right)^{-\frac{2m}{k}} \rightarrow 0 \quad \text{as } m \rightarrow \infty$$

for any $t_2 < t_3 < T$ and $a_2 < a_3 < a_\dagger$, and thus $w \equiv 0$ for $(t, a, x) \in Q_{T, a_\dagger}^\Omega$.

In case $\beta = 1$, \mathcal{I}_5 in (3.35) can be directly estimated from (3.37), using the Lipschitz continuity in (A_6) , without the need of employing the inequality (2.1) to obtain (3.38). The rest of details in obtaining $w \equiv 0$ in case $\beta = 1$ are skipped.

In conclusion, we can state the following uniqueness theorem.

Theorem 3.5. *Assume (A_1) - (A_6) hold. Then, the problem (1.4) satisfying the Robin boundary condition (1.6) admits no more than one solution in $\widetilde{W}_{T, a_\dagger}^\Omega$.*

ACKNOWLEDGMENTS

This work is in commemoration of the first death anniversary of V.A.K's father. V.A.K thanks Prof. Nguyen Huy Tuan for introducing him the ultra-parabolic problem.

REFERENCES

- [1] A. Friedman. *Partial Differential Equations of Parabolic Type*. R.E. Krieger Publishing Company, 1983.
- [2] S.A. Gourley, R. Liu, and J. Wu. Some vector borne diseases with structured host populations: extinction and spatial spread. *SIAM Journal on Applied Mathematics*, 67(2):408–433, 2007.
- [3] C.O. Horgan. Eigenvalue estimates and the trace theorem. *Journal of Mathematical Analysis and Applications*, 69:231–242, 1979.
- [4] F.C. Jiang and F.W. Meng. Explicit bounds on some new nonlinear integral inequality with delay. *Journal of Computational and Applied Mathematics*, 205:479–486, 2007.
- [5] J.M. Lee. *Introduction to Smooth Manifolds*. Graduate Texts in Mathematics, Volume 218. Springer-Verlag, New York, 2003.

- [6] L. Lorenzi. An abstract ultraparabolic integrodifferential equation. *Matematiche*, 58(2):401–435, 1998.
- [7] L. Lorenzi. An identification problem for an ultraparabolic integrodifferential equation. *Journal of Mathematical Analysis and Applications*, 234:417–456, 1999.
- [8] P. Magal and S. Ruan. *Structured Population Models in Biology and Epidemiology*. Mathematical Biosciences Subseries, Volume 1936. Springer-Verlag, Berlin, 2008.
- [9] M.D. Marozzi. On the valuation of Asian options by variational methods. *SIAM Journal on Scientific Computing*, 24:1124–1140, 2003.
- [10] M.D. Marozzi. Extrapolation discontinuous Galerkin method for ultraparabolic equations. *Journal of Computational and Applied Mathematics*, 224:679–687, 2009.
- [11] A.G. McKendrick. Applications of mathematics to medical problems. *Proceedings of the Edinburgh Mathematical Society*, 44:98–130, 1926.
- [12] J.W.H. So, J. Wu, and X. Zou. A reaction diffusion model for a single species with age structure, I. Travelling wave fronts on unbounded domains. *Proceedings of the Royal Society A: Mathematical, Physical & Engineering Sciences*, 457:1841–1853, 2001.
- [13] H. von Foerster. *Some remarks on changing populations*, In: *The Kinetics of Cellular Proliferation*. 382–407. Grune & Stratton, New York, 1959.
- [14] M. Witten. Modeling cellular systems and aging processes. I. Mathematics of cell system models: A review. *Mechanisms of Ageing and Development*, 17:53–94, 1981.
- [15] F. Zouyed and F. Rebbani. A modified quasi-boundary value method for an ultraparabolic ill-posed problem. *Journal of Inverse and Ill-Posed Problems*, 22(4):449–466, 2014.

INSTITUTE FOR NUMERICAL AND APPLIED MATHEMATICS, UNIVERSITY OF GOETTINGEN, GERMANY

FACULTY OF SCIENCES, HASSELT UNIVERSITY, CAMPUS DIEPENBEEK, BELGIUM
E-mail address: `vakhua.hcmus@gmail.com`

FACULTY OF SOLID STATE PHYSICS, GDANSK UNIVERSITY OF TECHNOLOGY, POLAND
E-mail address: `thehung.tran@mathmods.eu`

DEPARTMENT OF APPLIED MATHEMATICS, UNIVERSITY OF LEEDS, UNITED KINGDOM
E-mail address: `amt51d@maths.leeds.ac.uk`



UHasselT Computational Mathematics Preprint Series

2018

- UP-18-07 *Vo Anh Khoa, Tran The Hung, Daniel Lesnic*, **Uniqueness result for an age-dependent reaction-diffusion problem**, 2018
- UP-18-06 *Koondanibha Mitra, Iuliu Sorin Pop*, **A modified L-Scheme to solve nonlinear diffusion problems**, 2018
- UP-18-05 *David Landa-Marban, Na Liu, Iuliu Sorin Pop, Kundan Kumar, Per Pettersson, Gunhild Bodtker, Tormod Skauge, Florin A. Radu*, **A pore-scale model for permeable biofilm: numerical simulations and laboratory experiments**, 2018
- UP-18-04 *Florian List, Kundan Kumar, Iuliu Sorin Pop and Florin A. Radu*, **Rigorous upscaling of unsaturated flow in fractured porous media**, 2018
- UP-18-03 *Koondanibha Mitra, Hans van Duijn*, **Wetting fronts in unsaturated porous media: the combined case of hysteresis and dynamic capillary**, 2018
- UP-18-02 *Xiulei Cao, Koondanibha Mitra*, **Error estimates for a mixed finite element discretization of a two-phase porous media flow model with dynamic capillarity**, 2018
- UP-18-01 *Klaus Kaiser, Jonas Zeifang, Jochen Schütz, Andrea Beck and Claus-Dieter Munz*, **Comparison of different splitting techniques for the isentropic Euler equations**, 2018

2017

- UP-17-12 *Carina Bringedal, Tor Eldevik, Øystein Skagseth and Michael A. Spall*, **Structure and forcing of observed exchanges across the Greenland-Scotland Ridge**, 2017

- UP-17-11 *Jakub Wiktor Both, Kundan Kumar, Jan Martin Nordbotten, Iuliu Sorin Pop and Florin Adrian Radu, **Linear iterative schemes for doubly degenerate parabolic equations**, 2017*
- UP-17-10 *Carina Bringedal and Kundan Kumar, **Effective behavior near clogging in upscaled equations for non-isothermal reactive porous media flow**, 2017*
- UP-17-09 *Alexander Jaust, Balthasar Reuter, Vadym Aizinger, Jochen Schütz and Peter Knabner, **FESTUNG: A MATLAB / GNU Octave toolbox for the discontinuous Galerkin method. Part III: Hybridized discontinuous Galerkin (HDG) formulation**, 2017*
- UP-17-08 *David Seus, Koondanibha Mitra, Iuliu Sorin Pop, Florin Adrian Radu and Christian Rohde, **A linear domain decomposition method for partially saturated flow in porous media**, 2017*
- UP-17-07 *Klaus Kaiser and Jochen Schütz, **Asymptotic Error Analysis of an IMEX Runge-Kutta method**, 2017*
- UP-17-06 *Hans van Duijn, Koondanibha Mitra and Iuliu Sorin Pop, **Traveling wave solutions for the Richards equation incorporating non-equilibrium effects in the capillarity pressure**, 2017*
- UP-17-05 *Hans van Duijn and Koondanibha Mitra, **Hysteresis and Horizontal Redistribution in Porous Media**, 2017*
- UP-17-04 *Jonas Zeifang, Klaus Kaiser, Andrea Beck, Jochen Schütz and Claus-Dieter Munz, **Efficient high-order discontinuous Galerkin computations of low Mach number flows**, 2017*
- UP-17-03 *Maikel Bosschaert, Sebastiaan Janssens and Yuri Kuznetsov, **Switching to nonhyperbolic cycles from codim-2 bifurcations of equilibria in DDEs**, 2017*
- UP-17-02 *Jochen Schütz, David C. Seal and Alexander Jaust, **Implicit multiderivative collocation solvers for linear partial differential equations with discontinuous Galerkin spatial discretizations**, 2017*
- UP-17-01 *Alexander Jaust and Jochen Schütz, **General linear methods for time-dependent PDEs**, 2017*

2016

- UP-16-06 *Klaus Kaiser and Jochen Schütz, **A high-order method for weakly compressible flows**, 2016*

- UP-16-05 *Stefan Karpinski, Iuliu Sorin Pop, Florin A. Radu*, **A hierarchical scale separation approach for the hybridized discontinuous Galerkin method**, 2016
- UP-16-04 *Florin A. Radu, Kundan Kumar, Jan Martin Nordbotten, Iuliu Sorin Pop*, **Analysis of a linearization scheme for an interior penalty discontinuous Galerkin method for two phase flow in porous media with dynamic capillarity effects** , 2016
- UP-16-03 *Sergey Alyaev, Eirik Keilegavlen, Jan Martin Nordbotten, Iuliu Sorin Pop*, **Fractal structures in freezing brine**, 2016
- UP-16-02 *Klaus Kaiser, Jochen Schütz, Ruth Schöbel and Sebastian Noelle*, **A new stable splitting for the isentropic Euler equations**, 2016
- UP-16-01 *Jochen Schütz and Vadym Aizinger*, **A hierarchical scale separation approach for the hybridized discontinuous Galerkin method**, 2016

All rights reserved.